

## Antibiosis and GC-MS of secondary metabolites of rhizosphere bacteria from Manatee food plants in the humic freshwater ecosystem of Eniong river, Nigeria

J. Ekong<sup>a</sup>, O. K. Fatunla<sup>a,\*</sup>, B. N. Ita<sup>b</sup>, N. U. Asamudo<sup>a</sup>, O. U. M. John<sup>a</sup>, N. A. Abraham<sup>a</sup>, J. P. Essien<sup>a</sup>

<sup>a</sup>Department of Microbiology, University of Uyo, Akwa Ibom state, Nigeria

<sup>b</sup>Department of Chemistry, University of Uyo, Akwa Ibom state, Nigeria

### ARTICLE INFO

#### Article history:

Received: 12 January 2024

Received in revised form: 05 May 2024

Accepted: 24 May 2024

Available online: 12 June 2024

**Keywords:** Manatee, Food Plants, Antibiosis, Bioactive Compounds, Antibiosis

DOI:10.61298/rans.2024.2.1.68

### ABSTRACT

Microorganisms are able to synthesize secondary metabolites of various structures and bioactivities. These metabolites are produced to help the organism compete successfully with other organisms in their natural habitat and adapt with changing environmental milieu. The ability of rhizosphere bacteria (*Bacillus subtilis* NC\_000964.3 and *Pseudomonas aeruginosa* NC\_002516.2) isolated from the rhizospheric soil of Manatee food plants *Mimosa pygra*, *Ipomeoa aquatica* and *Pistia stratoites* to inhibit the growth of human pathogens (*P. aeruginosa*, *E. coli*, *S. aureus* and *B. subtilis*) was evaluated using standard methods. It was observed that the growth extracts of *B. subtilis* strains M<sub>5</sub>, M<sub>8</sub> and P<sub>7</sub> and *P. aeruginosa* strains I<sub>3</sub> and M<sub>9</sub> contained useful bioactive compound. GC-MS analysis of the cell-free methanol extract of the antibiotic producing bacterial strains was also evaluated and the results showed that their inhibitory potentials against bacterial pathogens are due to the presence of phenylethyl alcohol, 2-ethyl-4-methyl-1,3-dioxolane, bicyclo [4.2.0] octa-1,3,5-triene and 4-amino-2-methyl-5,6-dimethyl pyrimidine for *B. subtilis* and 3,4-dimethyl tetrahydrofuran, 4,6-dimethyl-4-hydroxy-5-heptenoic acid and 2,4-dimethyl-4-heptanol for *Pseudomonas aeruginosa*. These strains of rhizosphere bacteria may be exploited to produce new antibiotics.

©2024 The Author(s). Production and Hosting by FLAYOO Publishing House LTD on behalf of the Nigerian Society of Physical Sciences (NSPS). Peer review under the responsibility of NSPS. This is an open access article under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

### 1. INTRODUCTION

A phenomenon known as antibiosis occurs when the growth of one or more organisms is hampered by the presence of other species or the release of their metabolites. Antibiotics are chemical or bioactive metabolites created by bacteria that, in small doses, have the power to prevent other organisms from growing, metabolizing, and even killing them. The role of natural antibiotics has been disputed for decades and has been the

subject of much investigation [1]. Natural antibiotics are hypothesized to aid in microbial defense, fitness, interference, and competition. Antibiotic-producing organisms are widespread in microbial communities around the world [1–3]. It is well acknowledged that in terrestrial ecosystems, naturally occurring antibiotics accumulate at inhibiting concentrations largely in nutrient-rich environments, including those that antibiotic producers come across when colonizing the rhizosphere of plants or animal hosts [1].

Numerous crucial environmental tasks are carried out by the freshwater ecosystem, including nutrient recycling, water purifi-

\*Corresponding author: Tel.: +234-806-4049-745.

e-mail: opeyemifatunla@uniuyo.edu.ng (O. K. Fatunla)

cation, flood mitigation, groundwater recharge, and the provision of habitat for species. Additionally, people utilize them for recreation, especially in coastal areas [4]. This environment supports a wide variety of hydrophytes, vertebrate and non-vertebrate species, and microbiomes because to its high biological diversity. *Pseudomonas*, *Bacillus*, *Azotobacter*, *Micrococcus*, *Enterococcus*, *Acromobacterium*, *Salmonella*, *Shigella*, *Citrobacter*, *Flavobacterium*, and *Escherichia species* are only a few of the known bacterial species. There are additional known instances of fungi from the genera *Penicillium*, *Aspergillus*, *Candida*, *Fusarium*, *Geotricum*, *Saccharomyces*, and others [5]. These microbes are crucial to the mineralization of the intricate organics and other substances found in freshwater sediment. Within plants rhizosphere, considerable microbial activity occurs. The microbes present modulate biogeochemical processes within this system [6]. Simulation of bacteria is due to release from plant roots of a vast majority of plant materials including carbohydrates, vitamins, amino acids, enzymes, mucilage, sloughed root cells and carbon allocated to root-associated symbionts [7, 8]. This in turn affects the number and diversity of microorganisms present within the rhizosphere, their composition and interactions [9].

Recent studies have expanded our understanding of rhizosphere bacteria in aquatic systems, highlighting their unique roles and interactions within their environments. Findings by Pan *et al.* [10] & Lu *et al.* [11] demonstrated that aquatic rhizosphere bacteria, similar to their terrestrial counterparts, produce a diverse array of secondary metabolites that influence nutrient cycling and plant health. These findings underscore the complexity and dynamism of microbial interactions in semi-aquatic environments, where water flow and sediment composition add additional layers of ecological interaction not commonly observed in terrestrial habitats.

Furthermore, comparative analyses of secondary metabolite production in terrestrial and aquatic rhizospheres have revealed significant differences in the chemical profiles of produced antibiotics, likely due to the distinct selective pressures exerted by the aquatic environment [12, 13]. These studies suggest that aquatic rhizosphere bacteria might employ unique biochemical pathways to synthesize secondary metabolites, which could offer novel bioactive compounds for medical and agricultural applications.

The impact of these metabolites on microbial community dynamics and plant interactions in aquatic systems has also been explored in recent research. For example, a study by Etesami & Glick [14] in coastal marshlands showed that secondary metabolites from rhizosphere bacteria significantly affect the growth patterns of both plant hosts and neighboring microbial populations, potentially influencing the overall ecosystem stability and resilience to environmental stressors.

Moreover, the exploration of antibacterial activity in these environments has gained momentum, with studies such as those by Saeed *et al.* [15] identifying potent antimicrobial agents from bacteria isolated from the rhizosphere of aquatic plants. These agents not only offer potential for therapeutic uses but also play crucial ecological roles, mediating interactions within the microbial community and protecting plant hosts against pathogens.

Rhizosphere bacteria have been documented to serve as biocontrol and antibacterial agents in addition to influencing plant

development. Raaijmakers [16], Weller *et al.* [17], and Mavrodi *et al.* [18] all found that rhizosphere-dwelling biocontrol bacteria may suppress plant diseases through the use of antibiotics. Examples of antibiosis that protects eukaryotic hosts and their symbionts from pathogens include associations between ants and *Pseudonocardia* species, which produce antibiotics that shield ant fungal gardens from the mycoparasite *Escovopsis* [19], and the southern pine beetle *Dendroctonus frontalis* and the fungus *Entocorticium* sp. A, which is shielded [20]. The genera *Pseudomonas* and *Bacillus* isolated from the rhizosphere of maize and potatoes plant showed antagonistic activity against *Aspergillus flavus*, *Fusarium verticillioides* and phytopathogenic fungi such as *Fusarium osyosporum*, *Fusarium solani*, *Rhizoctonia solani* and *Pythium ultimum* respectively [20, 21]. Rhizosphere bacteria isolated from Kochi, India, showed positive activity against *Enterococcus* sp and *Staphylococcus aureus* [22]. Also, Nair *et al.* [23] isolated antibacterial compound from *Bacillus horikoshii* found in the rhizosphere region of Alfalfa plant. In addition, it has been observed that the growth of *Micrococcus luteus* and *Staphylococcus aureus* is inhibited by other soil bacteria, including *Streptomyces* species, *Pseudomonas aeruginosa*, *Actinomyces* strain, and *Bacillus* species [24, 25]. These findings suggest that soil microbes are good sources of secondary metabolites with potent antimicrobial properties.

Mammals get their gut bacteria mostly from the food they consume [26]. Recent research has revealed that metabolites isolated from commensal microorganisms of aquatic mammals possess a variety of antimicrobial activities. Studies on faecal specimens from terrestrial mammalian species have shown striking degrees of host specificity of microbiota, reflecting the influence of host phylogeny, gut anatomy, and diet [26]. It is well recognized that certain of these metabolites alter how the host behaves, which may have an impact on a person's susceptibility to illness and pathogen invasion. The West African manatee, or *Trichechus senegalensis*, is a big aquatic animal that is also known as a sea cow. *Ipomoea aquatica*, *Echinochloa stagnina*, *Echinochloa stagnina*, *Nymphaea lotus*, *Echinochloa crusgavonis*, *Pistia stratiotes*, *Vossia cuspidate*, *Mimosa pygra*, and *Nymphaea lotus* are noTable examples of the aquatic and emergent plants that they consume. Manatees have recently been wrongfully and illegally taken from the humic freshwater habitat of the Eniong River in Nigeria's Akwa Ibom State. When the animals were killed, the intestines were dominated by newly cropped specimens of the plants *Mimosa pygra*, *Ipomoea aquatica*, and *Pistia stratiotes*. Recent arguments contend that antibiotics serve less as defensive agents and more as signaling molecules crucial to bacterial physiology, communication, and gene regulation. Antibiotics are thought to seldom accumulate in natural environments in inhibitory quantities [27]. According to Mavrodi *et al.* [18], antibiotics are likely to serve as signaling chemicals as well as antagonists in natural environments. Our understanding of the function played by these substances in nature, however, has been significantly hampered by our knowledge of the quantities, spatial and temporal patterns of antibiotic synthesis in the field. Natural antibiotics are notoriously difficult to detect and measure *in situ* because of microbial destruction, chemical breakdown, and/or binding to soil and organic materials, according to Bonsall [28] and Mavrodi *et al.* [18].

Although the mechanisms that affect the form and function of the microbiotas of terrestrial and aquatic animals are becoming more understood, relatively little is known about the factors that affect aquatic species that live in humic habitats. There are increasing concerns about the rising levels of antibiotic resistance, high cost and toxicity, the need to look for new therapeutic agents from soil microbiome has been on the increase. This is because soil microbiomes represent a promising habitat for the discovery and isolation of highly effective antibiotics as less than 1% of soil bacterial species are currently known. Information on the antibiotic potentials of bacterial isolates from the Niger Delta region of Nigeria is scanty, this prompted our search for new antibiotics from the rhizosphere of manatee food plants. Therefore, we report in this study the antibacterial activity and GC-MS analysis of methanolic extracts of bacteria from the rhizosphere of manatee food plants.

## 2. MATERIALS AND METHODS

The research area is a humic environment of the Eniong River, a tributary of the Cross River's middle course, and is situated on Nigeria's southern coast in the Niger Delta. The river lies between latitude 5°12'N – 5°22'N and longitude 7°54'E – 8°20'E. Rhizosphere samples (plant root and associated sediment) of three commonly encountered Manatee food plants (*Mimosa pygra* (Figure 1a), *Ipomeoa aquatica* (Figure 1b) and *Pistia stratiotes* (Figure 1c) detected in the stomach of dead manatee were collected manually from 10 different locations within the study area. All rhizosphere samples collected were placed in sterile tubes, transported to the laboratory and stored at 4°C for no more than 24 h before processing. Due to the presence of humic chemicals and soluble iron complexes, the river has a vivid coloration. Additionally, it supports a variety of aquatic animal species, including the critically endangered manatee (*Trichechus senegalensis* (Figure 2). Isolation of bacteria was carried out using serial dilution technique as described by Willey *et al.* [29]. Bacterial isolates were obtained on nutrient agar slant (NA) using the pour plate technique [30].

A sensitivity assay was carried out to determine the antibiotic producing ability of the bacterial isolates against pure and clinical isolates of *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Escherichia coli* and *Bacillus subtilis* using the crowded technique. Assay plates were incubated for 24hrs at 37°C. The method of Maniatis *et al.* [31] and Kraft *et al.* [32] were employed to extract and cure the plasmid from the antibiotic – producing isolates. Preliminary identification of the antibiotic producing bacterial strain revealed a Gram positive, cocci in pairs (EHS<sub>A4</sub>) as the best producer [33, 34]. Confirmation was done by the 16S rRNA gene sequence analysis. DNA was extracted from the pure culture by a salting –out procedure [35] and amplified by polymerase chain reaction using primers and PCR conditions adapted from Tuleva *et al.* [36] with forward primer 12F1 (5'CGTGCTTAACACATGCAA3') and the reverse primer 1390 (5'GCCACCGGCTTCGGGTGTTTA 3'). The amplified products were electrophoresed using 2% agarose gel and stained with ethidium bromide. The Sanger sequencing approach was used to do DNA sequencing utilizing an automated PCR cycle - Sanger Sequencer<sup>TM</sup> 3730/3730XL DNA analyser (Applied Systems). Direct blasting on <http://blast.ncbi.nlm.nih.gov> yielded the nu-

cleotide sequence. Using the agar well diffusion technique, antibiotic susceptibility of clinical isolates of *Pseudomonas aeruginosa*, *Escherichia coli*, *Staphylococcus aureus*, and *Bacillus subtilis* to methanolic growth extracts of antibiotic manufacturers was assessed [37]. The zones of inhibition were measured and recorded. Ciprofloxacin (100µg/mL) was used as the positive control. Whereas, bioactive compounds present in potent antibiotic-producing bacterial isolates were characterized and identified in accordance with the method of Elleuch [37]. GC-MS was performed on the methanolic extract of the crude culture medium (3mg/0.2mL), which had passed through a 0.45µm syringe filter to remove bacterial cells. A Hewlett-Packard 5890 gas chromatograph coupled with an AutoSpec mass spectrometer was used. Identification of individual compounds was done by comparison with the mass spectra of authentic reference compounds and reference libraries.

## 3. RESULTS

Thirteen (13) bacterial species including *Bacillus subtilis*, *Streptococcus* sp, *Proteus* sp, *Clostridium* sp, *Micrococcus* sp, *Serratia* sp, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Chromatium* sp, *Klebsiella* sp, *Flavobacterium* sp, *Actinomyetes* and *Enterobacter* sp were isolated from the rhizosphere of Manatee food plants. Their Gram staining characteristics is given in Table 1. Extracts from *B. subtilis* and *P. aeruginosa* were active against clinical isolates of *E. coli* and *S. aureus*, with zones of inhibition greater than 10mm (Table 2). Isolates M<sub>5</sub>, M<sub>8</sub> and P<sub>7</sub> were found to be strains of *B. subtilis*, while isolates I<sub>3</sub> and M<sub>9</sub> were strains of *P. aeruginosa* based on their genomic properties (Table 3). GC-MS analysis of methanol extracts from broth cultures of *B. subtilis* and *P. aeruginosa* strains revealed variations in compounds elaborated (Table 4). Nine compounds were detected in broth culture of isolates M<sub>5</sub>, M<sub>8</sub> and P<sub>7</sub> strains from *B. subtilis* while seven bioactive compounds were detected in isolates I<sub>3</sub> and M<sub>9</sub> from *P. aeruginosa*. The chromatograms of the bioactive compounds are given in Figures 3 and 4 while the elucidated structure of the compounds are presented in Figure 5.

## 4. DISCUSSION

Gram staining characteristics of the bacterial strains from the root zone of manatee food plants showed the presence of thirteen (13) isolates, of which ten (10) were rod shaped, two (2) cocci, and one (1) filamentous with many having the ability for spore formation. They were identified as *P. aeruginosa*, *E. coli*, *B. subtilis* and species of *Streptococcus*, *Clostridium*, *Micrococcus*, *Serratia*, *Chromatium*, *Klebsiella*, *Flavobacterium*, *Proteus*, *Actinomyetes*, and *Enterobacter*.

The ability of rhizosphere bacteria isolates to inhibit the growth of human pathogens (*P. aeruginosa*, *E. coli*, *S. aureus* and *B. subtilis*) was also evaluated. Methanol extracts of isolates from *B. subtilis*(P<sub>7</sub>) exhibited promising antibacterial activity against clinical isolates of *E. coli* and *S. aureus* (inhibition zone 10 - 14mm), but was bacteriostatic to *P. aeruginosa* (inhibition zone 6- 8mm). Similarly, the methanol extract of *P. aeruginosa* isolates (I<sub>3</sub> and M<sub>9</sub>) was most active against clinical strains of *E. coli* and *S. aureus* (inhibition zone = 10-14mm), but bacteriostatic against *B. subtilis* (inhibition zone <8mm). These results corroborate with report in literature. *Bacillus* species



**Figure 1.** a = *Mimosa pygra*; b = *Ipomeoa aquatica*; c = *Pistia stratoites*.

from terrestrial and marine environment are known to be good sources of antibiotic compounds [38–40]. Similarly, the genera *Pseudomonas* has been reported to show biological activity [40]. Also, Gislin *et al.* [22] reported the antibacterial activity of *B. amyloliquefaciens* from the rhizosphere and diverse cultivation at Kochi, India against *Enterococcus* sp and *S. aureus*. Nair *et al.* [23] reported antibacterial activity of *Bacillus horikoshii* isolated from the rhizosphere of Alfalfa plant against *Klebsiella* and *S. aureus*. Similar reports exist for *actinomycetes*, *streptomyces*, *Pseudomonads* and other genera from rhizospheric soils [41, 42]. Therefore, our results suggest that *P. aeruginosa* and *B. subtilis* from the rhizosphere of manatee food plants could serve as potent sources of novel antibiotic compounds.

Molecular analysis of the antibiotic producing bacterial iso-

lates was done using the 16S rRNA gene amplification and sequencing, followed by BLAST analysis using the mega blast tool of GenBank (<http://blast.ncbi.nlm.nih.gov>). This method is generally accepted as a better method of bacterial identification in comparison to other phenotypic methods [22]. Using this method, gene sequence comparison of P<sub>7</sub>, M<sub>5</sub> and M<sub>8</sub> demonstrated 99% similarities to *B. subtilis* NC\_000964.3. Similarly, sequence comparison of I<sub>3</sub> and M<sub>9</sub> demonstrated 99% similarities to *P. aeruginosa* NC\_002516.2, confirming the identity of the five bacterial strains (P<sub>7</sub>, M<sub>5</sub>, M<sub>8</sub>, I<sub>3</sub> and M<sub>9</sub>).

GC-MS analysis of the cell -free methanol extract of the antibiotic producing bacterial strains was also evaluated. *B. subtilis* produced a total of nine bioactive compounds from its culture broth. Similarly, *P. aeruginosa* synthesized seven

**Table 1. Gram staining characteristics of isolated bacterial species.**

Isolate code	Gram staining reaction	Shape	Probable organism
P <sub>7</sub> , M <sub>5</sub> , M <sub>8</sub>	+	Thick rod	<i>Bacillus subtilis</i>
P <sub>1</sub>	+	Cocci in chain	<i>Streptococcus</i> sp
I <sub>6</sub> , M <sub>1</sub>	-	Rod	<i>Proteus</i> sp
M <sub>6</sub>	+	Thick rod	<i>Clostridium</i> sp
M <sub>3</sub> , M <sub>7</sub>	+	Tiny cocci	<i>Micrococcus</i> sp
P <sub>4</sub> , M <sub>4</sub>	-	Rod	<i>Serratia</i> sp
P <sub>3</sub>	-	Short rod	<i>Escherichia coli</i>
I <sub>3</sub> , M <sub>9</sub>	-	Rod	<i>Pseudomonas aeruginosa</i>
P <sub>5</sub> , M <sub>4</sub>	-	Rod	<i>Chromatium</i> sp
I <sub>2</sub> , I <sub>4</sub>	-	Rod	<i>Klebsiella</i> sp
P <sub>2</sub> , P <sub>6</sub>	-	Rod	<i>Flavobacterium</i> sp
I <sub>1</sub>	+	Filamentous	<i>Actinomycetes</i>
I <sub>5</sub> , M <sub>2</sub>	-	Rod	<i>Enterobacter</i> sp

**Table 2. Gram staining characteristics of isolated bacterial species.**

Code	Isolates	Test clinical isolates			
		<i>P. aeruginosa</i>	<i>E. coli</i>	<i>S. aureus</i>	<i>B. subtilis</i>
P <sub>7</sub> M <sub>5</sub> M <sub>8</sub>	<i>B. subtilis</i>	+	-	-	+++
P <sub>1</sub>	<i>Streptococcus</i> sp	-	-	-	-
I <sub>6</sub> , M <sub>1</sub>	<i>Proteus</i> sp	-	-	-	-
M <sub>6</sub>	<i>Clostridium</i> sp	-	-	-	-
M <sub>3</sub> , M <sub>7</sub>	<i>Micrococcus</i> sp	-	-	-	-
P <sub>4</sub> , M <sub>4</sub>	<i>Serratia</i> sp	-	-	-	-
P <sub>3</sub>	<i>E. coli</i>	-	-	-	-
I <sub>3</sub> , M <sub>9</sub>	<i>P. aeruginosa</i>	-	+++	+++	+
P <sub>5</sub> , M <sub>4</sub>	<i>Chromatium</i> sp	-	-	-	-
I <sub>2</sub> , I <sub>4</sub>	<i>Klebsiella</i> sp	-	-	-	-
P <sub>2</sub> , P <sub>6</sub>	<i>Flavobacterium</i> sp	-	-	-	-
P <sub>1</sub>	<i>Actinomycetes</i>	-	-	-	-
I <sub>5</sub> , M <sub>2</sub>	<i>Enterobacter</i> sp	-	-	-	-

Key: 0-6mm = -, 6-8mm = +, 8-10mm = ++, 10-14mm = +++.

**Table 3. Genomic properties of bioactive strains.**

Isolate number	Sample type	DNA type	Gene bank accession number	ID of organism
1	P <sub>7</sub>	DNA	NC_000964.3	<i>B. subtilis</i>
2	M <sub>8</sub>	DNA	NC_000964.3	<i>B. subtilis</i>
3	I <sub>3</sub>	DNA	NC_002516.2	<i>P. aeruginosa</i>
4	M <sub>9</sub>	DNA	NC_002516.2	<i>P. aeruginosa</i>
5	M <sub>5</sub>	DNA	NC_000964.3	<i>B. subtilis</i>

bioactive components. For extract of *B. subtilis* strains, 2-ethyl-2-pentanal (22.25%) and 2-ethyl-2-hexenal (21.49%) were the major compounds detected. Others were 2-ethyl-4-methyl-1,3-dioxalane (17.91%), 3-octanol (11.91%), trimethyl silanol (7.25%), 4-amino-2-methyl-5,6-dimethyl pyrimidine (5.77%), cyclohexane carboxylic acid (4.97%), bicyclo [4.2.0] octa-1,3,5-triene (4.51%) and 4-methyl-1,3-dioxane (3.95%). For *P. aeruginosa* extract, the major compound detected was 4-methyl-4-octanol (35.31%), followed by 2-ethyl-2-hexenal (16.77%), 2-ethyl-2-pentanal (15.05%), 2-methyl-2-hexanol (12.67%), 1,1-dimethoxy-2-methylpropane (8.26%), 2-methoxy quinoline (6.57%), and isothiocyanato ethane (5.37%). The antimicrobial activity of phenolic compounds and aromatic alcohols such as



phenylethyl alcohol, have been reported [43]. Their bactericidal action was related to their physicochemical properties, and involved alterations in membrane function, resulting in cell death, particularly for Gram – negative bacteria. The antibacterial activity of fatty alcohols has been documented. Alcohols with carbon length between 8 and 12 showed activity against *S. aureus* and *P. acnes* [44]. These alcohols disrupt cell membrane fluidity by infiltrating the membrane's molecular structure, with the polar hydroxyl group hydrogen-bonded into the aqueous phase and the nonpolar carbon chain aligned into the lipid phase by dispersion forces. The antibacterial strains' extracts included cyclohexane carboxaldehyde, 2-ethyl-2-pentanal, and 2-ethyl-2-hexenal. Antibacterial activity has been observed for aliphatic

**Table 4. Bioactive compounds from methanol extract of broth cultures from rhizosphere bacterial isolates.**


Isolate	RT	Area (%)	Compound
<i>B. subtilis</i> strains M <sub>5</sub> , M <sub>8</sub> and P <sub>7</sub> .	3.297	4.51	Bicyclo [4.2.0] octa-1,3,5-triene
	3.387	4.97	Cyclohexane carboxaldehyde
	3.600	22.25	2-ethyl-2-pentenal
	3.657	11.91	3-octanol
	3.773	17.91	2-ethyl-4-methyl-1,3-dioxolane
	4.144	21.49	2-ethyl-2-hexenal
	4.271	3.95	4-methyl-1,3dioxane
	4.942	7.25	Trimethyl silanol
	6.667	5.77	4-amino-2-methyl-5,6-dimethyl pyrimidine
<i>P. aeruginosa</i> strains I <sub>3</sub> and M <sub>9</sub> .	3.660	15.05	2-ethyl-2-pentenal
	3.709	12.67	2-methyl-2-hexanol
	3.818	5.37	Isothiocyanato-ethane
	4.178	16.77	2-ethyl-2-hexenal
	4.369	35.31	4-methyl-4-octanol
	4.950	8.26	1,1-dimethoxy-2-methyl-propane
	5.880	6.57	2-methoxy-quinoline

**Figure 2.** Manatee (*Trichechus senegalensis*) encountered during the study.

aldehydes [45]. These aldehydes work by changing the function of membrane-associated proteins, interacting with the cell membrane's nucleophilic groups and producing considerable disruption in the lipidic bilayer. This action is more noticeable for the  $\alpha$ ,  $\beta$  unsaturated aldehydes. According to Kim *et al.* [46], silanols are a novel family of antibacterial drugs that appear to be more effective than their comparable alcohols. The authors examined the antibacterial activities of silanols, alcohols, and phenols on *E. coli* C3000 (ATCC15597), a laboratory strain of *S. aureus*, *P. aeruginosa*, and *E. faecalis* and discovered that silanols with stronger antibacterial activity had lower MIC values. Quinolones, on the other hand, have been shown to exhibit

**Figure 3.** Profile of bioactive compounds elaborated by *B. subtilis* strains M<sub>5</sub>, M<sub>8</sub> and P<sub>7</sub>.**Figure 4.** Profile of bioactive compounds elaborated by *Pseudomonas aeruginosa* strains I<sub>3</sub> and M<sub>9</sub>.

a wide range of biological actions, including antibacterial [47], antimalarial [48], anti-inflammatory [49], cytotoxicity [50], and so on. These compounds may be responsible for the extracts' antibacterial effect, either alone or in combination. In contrast to other species, the bioactive *B. subtilis* and *P. aeruginosa* strains



**Figure 5.** Bioactive compounds from methanolic extracts of A (*Bacillus subtilis* NC\_000964.3) and the bioactive compounds from methanolic extracts B (*Pseudomonas aeruginosa* NC\_002516.2).

support previous findings. Gislin *et al.* [22] discovered that *Bacillus* species are rich in antibiotic substances. This is consistent with the findings of this study. Sharma and Kaur [51] investigated the antibacterial activity of *Pseudomonas* and *Bacillus* rhizobacterial strains obtained from carnation rhizosphere soil. *Bacillus* sp isolated from the Amazon basin has also been shown to have antibacterial properties [52]. Iqbal *et al.* [53] identified antibacterial peptides from *B. safensis* strain MK12 waste dump soil in Pakistan and screened, characterized, and optimized them.

The bioactive compounds identified in our study, derived from the rhizosphere bacteria of manatee food plants, show promising antibacterial activity. To understand the potential mechanisms through which these compounds exert their effects, it is crucial to consider their molecular interactions with bacterial cells. This understanding is pivotal for leveraging their therapeutic potential and ecological significance. The primary mode of action of many natural bioactive compounds involves disrupting the bacterial cell membrane integrity. This disruption can lead to increased cell permeability, resulting in the leakage of vital cellular contents and eventual cell death. For instance, compounds such as lipopeptides and polyketides, commonly produced by soil bacteria, are known to insert themselves into the lipid bilayer of bacterial membranes, distorting the membrane structure and causing functional disruptions [54, 55].

Another potential mechanism is the inhibition of protein synthesis. Some bioactive compounds bind to bacterial ribosomes, blocking the ribosomal tunnel used during protein translation. This action effectively halts the protein production necessary for bacterial growth and replication, a mechanism employed by several known antibiotics like tetracyclines and macrolides [56]. Given the diverse microbial origin of the compounds studied, similar interactions could be involved in their antibacterial activity. Furthermore, these compounds might interfere with the synthesis or function of essential bacterial enzymes. Enzymatic inhibition can occur through the binding of bioactive compounds to the active sites of key bacterial enzymes, altering their conformation and thereby inhibiting their catalytic activity. This type of inhibition is crucial as it affects bacterial metabolic pathways and can lead to the buildup of toxic intermediates or the depletion of essential metabolites necessary for cell survival [57]. Bioactive compounds may also affect bacterial DNA replication. Certain

compounds are capable of intercalating into DNA or binding to DNA gyrase, an enzyme critical for DNA replication. By disrupting these processes, the compounds prevent the bacteria from reproducing and maintaining their genomic integrity [58].

## 5. CONCLUSION

Secondary metabolites with a range of structures and bioactivities can be produced by microorganisms. Nutrients, growth rate, enzyme induction, and enzyme inactivation all affect how they are produced. The production of these metabolites, which have been crucial in the discovery and development of several antibiotics, helps the organism compete successfully with other species in its natural habitat and adapt to changing environmental conditions. As a result, a significant portion of commercially available antimicrobial medicines have microbial origins. The results of the current investigation have demonstrated that bacteria from the rhizosphere of manatee food plants can be powerful sources of new antibiotics and other bioactive substances.

## ACKNOWLEDGEMENTS

Authors express appreciation to International Centre for Energy and Environmental Sustainability Research (ICEESR), Nigeria, for providing partial support of this study.

Partial financial support was received from International Centre for Energy and Environmental Sustainability Research (ICEESR), Nigeria to cover the cost of some analysis.

## References

- [1] D. V. Mavrodi, W. Blankenfeldt & L. S. Thomashow, "Phenazine compounds in fluorescent *Pseudomonas* spp.: biosynthesis and regulation", *Annu. Rev. Phytopathol* **44** (2006) 417. <https://doi.org/10.1146/annurev.phyto.44.013106.145710>.
- [2] B. M. Wiseman, S. M. Neate, K. O. Keller & S. E. Smith, "Suppression of *Rhizoctonia solani* anastomosis group 8 in Australia and its biological nature", *Soil Biol Biochem* **28** (1996) 727. [https://doi.org/10.1016/0038-0717\(95\)00178-6](https://doi.org/10.1016/0038-0717(95)00178-6).
- [3] J. J. Scott, "Bacterial protection of beetle-fungus mutualism", *Science* **322** (2008) 63. <https://pubmed.ncbi.nlm.nih.gov/18832638/>.
- [4] G. R. Daily, "Nature's services: societal dependence on natural ecosystems", *Environmental Values* **7** (1998) 365. <https://philpapers.org/rec/DAINSS-2>.
- [5] P. A. Del Giorgio & J. J. Cole, "Bacterial energetics and growth efficiency", in *Microbial ecology of the oceans*, Wiley-Liss, 2000, pp.

- 289–325. [https://www.researchgate.net/publication/262403297\\_Bacterial\\_energetics\\_and\\_growth\\_efficiency](https://www.researchgate.net/publication/262403297_Bacterial_energetics_and_growth_efficiency).
- [6] J. Li, G. Ren, Z. Jia & Y. Dong, "Composition and activity of rhizosphere microbial communities associated with healthy and diseased greenhouse tomatoes", *Plant Soil* **380** (2014) 337. <https://doi.org/10.1007/s11104-014-2097-6>.
- [7] P. G. Dennis, A. J. Miller & P. R. Hirsch, "Are root exudates more important than other sources of rhizodeposits in structuring rhizosphere bacterial communities?" *FEMS Microbiology Ecology* **72** (2010) 313. <https://doi.org/10.1111/j.1574-6941.2010.00860.x>.
- [8] M. Koranda, J. Schnecker, C. Kaiser, L. Fuchsluger, B. Kitzler, C. F. Stange, A. Sessitsch, S. Zechmeister-Boltenstern & A. Richter, "Microbial processes and community composition in the rhizosphere of European beech - The influence of plant C exudates", *Soil Biology and Biochemistry* **43** (2011) 551. <https://doi.org/10.1016/j.soilbio.2010.11.022>.
- [9] E. Somers, J. Vanderleyden & M. Srinivasan, "Rhizosphere bacterial signalling: A love parade beneath our feet", *Critical Review of Microbiology* **30** (2004) 205. <https://doi.org/10.1080/10408410490468786>.
- [10] X. Pan, J. M. Raaijmakers & V. J. Carrión, "Importance of Bacteroidetes in host-microbe interactions and ecosystem functioning", *Trends in Microbiology* **31** (2023) 842. <https://doi.org/10.1016/j.tim.2023.03.018>.
- [11] Y. Lu, H. J. Kronzucker, M. Yu, S. Shabala & W. Shi, "Nitrogen-loss and carbon-footprint reduction by plant-rhizosphere exudates", *Trends in Plant Science* **59** (2023) 959. <https://doi.org/10.1016/j.tplants.2023.09.007>.
- [12] A. Kumar & A. Dubey, "Rhizosphere microbiome: Engineering bacterial competitiveness for enhancing crop production", *Journal of Advanced Research* **24** (2020) 337. <https://doi.org/10.1016/j.jare.2020.04.014>.
- [13] V. Mathur & D. Ulanova, "Microbial metabolites beneficial to plant hosts across ecosystems", *Microbial Ecology* **86** (2023) 25. <https://doi.org/10.1007/s00248-022-02073-x>.
- [14] H. Etesami & B. R. Glick, "Bacterial indole-3-acetic acid: A key regulator for plant growth, plant-microbe interactions, and agricultural adaptive resilience", *Microbiological Research* **281** (2024) 127602. <https://doi.org/10.1016/j.micres.2024.127602>.
- [15] Q. Saeed, W. Xiukang, F. U. Haider, J. Kučerik, M. Z. Mumtaz, J. Holatko & A. Mustafa, "Rhizosphere bacteria in plant growth promotion, biocontrol, and bioremediation of contaminated sites: a comprehensive review of effects and mechanisms", *International Journal of Molecular Sciences* **22** (2021) 105. <https://doi.org/10.3390/ijms221910529>.
- [16] J. M. Raaijmakers, M. Vlami & J. T. de Souza, "Antibiotic production by bacterial biocontrol agents", *Antonie Van Leeuwenhoek* **81** (2002) 537. <https://doi.org/10.1023/A:1020501420831>.
- [17] D. M. Weller, B. B. Landa, O. V. Mavrodi, K. L. Schroeder, L. E. De La Fuente, S. B. Bankhead, R. A. Molar, R. F. Bonsall, D. V. Mavrodi & L. S. Thomashow, "Role of 2,4-diacetylphloroglucinol-producing fluorescent *Pseudomonas* spp. in the defense of plant roots", *Plant Biol.* **9** (2007) 4. <https://pubmed.ncbi.nlm.nih.gov/17058178/>.
- [18] D. V. Mavrodi, O. V. Mavrodi, J. A. Parejko, R. F. Bonsall, Y. S. Kwak, T. C. Paulitz, L. S. Thomashow & D. M. Weller, "Accumulation of the antibiotic phenazine-1-carboxylic acid in the rhizosphere of dryland cereals", *Appl Environ Microbiol* **78** (2012) 804. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3264129/>.
- [19] C. R. Currie, M. Poulsen, J. Mendenhall, J. J. Boomsma & J. Billen, "Co-evolved crypts and exocrine glands support mutualistic bacteria in fungus-growing ants", *Science* **311** (2006) 81. <https://pubmed.ncbi.nlm.nih.gov/16400148/>.
- [20] J. D. Palumbo, T. L. O'keeffe & H. K. Abbas, "Isolation of Maize Soil and Rhizosphere Bacteria with Antagonistic Activity against *Aspergillus flavus* and *Fusarium verticillioides*", *Journal of Food Protection* **70** (2007) 1615. <https://pubmed.ncbi.nlm.nih.gov/16400148/>.
- [21] S. Mezaache-Aichour, A. Guechi, J. Nicklin, D. Drider, H. Prevost & R. N. Strange, "Isolation, identification and antimicrobial activity of pseudomonads isolated from the rhizosphere of potatoes growing in Algeria", *Journal of Plant Pathology* **94** (2012) 89. <https://hal.inrae.fr/view/index/identifiant/hal-02645667>.
- [22] D. Gislin, D. Sudarsanam, G. A. Raj & K. Baskar, "Antibacterial activity of soil bacteria isolated from Kochi, India and their molecular identification", *Journal of Genetic Engineering and Biotechnology* **16** (2018) 287. <https://doi.org/10.1016/j.jgeb.2018.05.010>.
- [23] N. M. Nair, R. Kanthasamy, R. Mahesh, S. I. K. Selvam & S. Ramalakshmi, "Identification of antibacterial compound from *Bacillus horikoshii*, isolated from Rhizosphere region of Alfalfa plant", *Journal of Applied Sciences* **19** (2019) 140. [https://ui.adsabs.harvard.edu/link\\_gateway/2019JApSc..19..140M/doi:10.3923/jas.2019.140.147](https://ui.adsabs.harvard.edu/link_gateway/2019JApSc..19..140M/doi:10.3923/jas.2019.140.147).
- [24] J. O. Falkinham, T. E. Wall, J. R. Tanner, K. Tawaha, F. Q. Alali, C. Li & N. H. Oberlies, "Proliferation of antibiotic-producing bacteria and concomitant antibiotic production as the basis for the antibiotic activity of Jordan's red soils", *Applied and Environmental Microbiology* **75** (2009) 2735. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2681674/>.
- [25] S. A. Hassan, E. Hanif & R. R. Zohra, "Isolation and screening of soil bacteria for potential antimicrobial activity", *FUUAST Journal of Biology* **4** (2014) 217. <https://fuuastjb.org/index.php/fuuastjb/article/view/172>.
- [26] R. E. Ley, M. Hamady, C. Lozupone, P. J. Turnbaugh, R. R. Ramey, J. S. Bircher, M. L. Schlegel, T. A. Tucker, M. D. Schrenzel, R. Knight & J. I. Gordon, "Evolution of Mammals and their Gut Microbes", *Science* **320** (2008) 1647. <https://pubmed.ncbi.nlm.nih.gov/18497261/>.
- [27] J. L. Martinez, "Antibiotics and antibiotic resistance genes in natural environments", *Science* **321** (2008) 365. <https://pubmed.ncbi.nlm.nih.gov/18635792/>.
- [28] R. F. Bonsall, L. S. Thomashow, D. V. Mavrodi & D. M. Weller, "Extraction and detection of antibiotics in the rhizosphere metabolome", *Curr. Trends Mass Spectrom. (Suppl. Spectrosc.)* **11** (2007) 14. <https://www.spectroscopyonline.com/view/extraction-and-detection-antibiotics-rhizosphere-metabolome>.
- [29] J. M. Willey, L. M. Sherwood & C. J. Woollerton, *Prescott Harley and Kleins Microbiology*, (7th Ed). Mc-Graw Hill, New York, 2008, pp. 1049–1088. <https://search.worldcat.org/title/Prescott-Harley-and-Klein's-microbiology/oclc/71044581>.
- [30] C. I. Ozoigwe & O. K. Agwa, "Microbiological quality of water collected from boreholes sited near refuse dumpsites in Port Harcourt, Nigeria", *African Journal of Biotechnology* **11** (2012) 3135. <https://www.ajol.info/index.php/ajb/article/view/100707>.
- [31] T. Maniatis, E. T. Fritsch & J. Sambrook, *Molecular Cloning, a laboratory manual*, Cold Spring Harbor: New York, 1989, pp. 945–982. <https://onlinelibrary.wiley.com/doi/abs/10.1016/0307-4412%2883%2990068-7>.
- [32] R. Kraft, J. Tardiff, K. S. Krauter & L. A. Leinwand, "Using mini-prep plasmid DNA for sequencing double stranded template with sequences", *Bio. Technique* **6** (1988) 544. <https://pubmed.ncbi.nlm.nih.gov/3273187/>.
- [33] D. J. Brenner, N. R. Krieg, J. T. Staley & G. M. Garrity, *Bergey's manual of systematic bacteriology, 2nd Edition Part B*, Springer's, USA, 2005, pp. 27–32. <https://link.springer.com/book/10.1007/0-387-28021-9>.
- [34] M. Cheesbrough, *District laboratory practices in tropical countries Second edition*, Cambridge University Press, 2006, pp. 13–23. <https://www.cambridge.org/core/books/district-laboratory-practice-in-tropical-countries/777AAC879D604058E7EA3F4E15824F88>.
- [35] S. A. Miller, D. D. Dykes & H. F. Polesky, "A simple salting out procedure for extracting DNA from human nucleated cells", *Nucleic Acids Res.* **16** (1988) 1215. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC334765/>.
- [36] B. Tuleva, N. Christova, R. Cohen, D. Antonova, T. Todrov & I. Stoineva, "Isolation and characterization of trehalose-tetraester biosurfactants from a soil strain *Micrococcus luteus*", *Process Biochemistry* **44** (2009) 135. <https://doi.org/10.1016/j.procbio.2008.09.016>.
- [37] L. Elleuch, M. Shaaban & S. Smaoui, "Bioactive secondary metabolites from a new terrestrial *Streptomyces* sp. TN262", *Appl. Biochem. Biotechnol* **162** (2010) 579. <https://doi.org/10.1007/s12010-009-8808-4>.
- [38] M. Kanagasabhapathy, H. Sasaki & S. Nagata, "Phylogenetic identification of epibiotic bacteria possessing antimicrobial activities isolated from red algal species of Japan", *World Journal of Microbiology and Biotechnology* **24** (2008) 2315. <https://doi.org/10.1007/s11274-008-9746-y>.
- [39] P. Devi, S. Wahidullah, C. Rodrigues & L. D. Souza, "The sponge-associated bacterium *Bacillus licheniformis* SAB1: A source of antimicrobial compounds", *Marine Drugs* **8** (2010) 1203. <https://doi.org/10.3390/md8041203>.
- [40] A. P. Graca, J. Bondoso, H. Gasper, J. R. Xavier, M. C. Monteiro, M. de la Cruz, D. Oves-Costales, F. Vincente & O. M. Lage, "Antimicrobial activity of heterotrophic bacterial communities from the marine sponge *Erylus discophorus* (Astrophorida, Geodiidae)", *PLOS ONE* **8** (2013) 1. <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0078992>.
- [41] J. Ramakrishnan, M. Shunmugasundaram & M. Narayanan, "Streptomyces sp. SCBT isolated from rhizosphere soil of medicinal plants is antagonistic to pathogenic bacteria", *Iran Journal of Biotechnology* **7** (2009) 75. <https://www.cabidigitallibrary.org/doi/full/10.5555/20093232987>.
- [42] M. F. Adegboye & O. O. Babalola, "Isolation and identification of potential antibiotic producing rare actinomycetes from rhizospheric soils", *Journal of Human Ecology* **56** (2016) 31. <https://doi.org/10.1080/09709274.2016>.

- 11907035.
- [43] J. J. Lucchini, J. Corre & A. Cremieux, "Antibacterial activity of phenolic compounds and aromatic alcohols", *Research in Microbiology* **141** (1990) 499. <https://www.sciencedirect.com/science/article/pii/S0923250890900752>.
- [44] I. Kubo, H. Muroi, H. Himejima & A. Kubo, "Antibacterial activity of long-chain alcohols: the role of hydrophobic alkyl groups", *Bioorg. Med. Chem. Lett.* **3** (1993) 1305. [https://doi.org/10.1016/S0960-894X\(00\)80336-4](https://doi.org/10.1016/S0960-894X(00)80336-4).
- [45] G. Bisignano, M. G. Lagana, D. Trombetta, S. Arena, A. Nostro, N. Uccella, G. Mazzanti & A. Saija, "In vitro antibacterial activity of some aliphatic aldehydes from *Olea europaea* L", *FEMS Microbiology Letters* **198** (2001) 9. <https://doi.org/10.1111/j.1574-6968.2001.tb10611.x>.
- [46] Y. Kim, S. Farrah & R. Baney, "Silanol—A novel Class of Antimicrobial Agent," *Electron J Biotech.* **9** (2006) 0. [https://www.scielo.cl/scielo.php?pid=S0717-34582006000200012&script=sci\\_arttext&tlng=en](https://www.scielo.cl/scielo.php?pid=S0717-34582006000200012&script=sci_arttext&tlng=en)
- [47] J. Eswaran, E. Koronakis, M. K. Higgins, C. Hughes & V. Koronakis, "Three's company: component structures bring a closer view of tripartite drug efflux pumps", *Curr Opin Struct Biol* **14** (2004) 741. <https://www.sciencedirect.com/science/article/abs/pii/S0959440X04001794>.
- [48] S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K. Averyt & M. Tignor, "Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change", Cambridge, UK: Cambridge University Press (2007). <https://wedocs.unep.org/handle/20.500.11822/30763;jsessionid=701380F0A26B27E3D4AB318AE8C63D77>.
- [49] P. Alam, S. I. Alqasoumi, F. Shakeel & M. S. Abdel-Kader, "HPTLC densitometric analysis of arbutin in bulk drug and methanolic extracts of *Arctostaphylos uva-ursi*", *Natural Product Research* **25** (2011) 1671. <https://www.tandfonline.com/doi/abs/10.1080/14786419.2010.529447>.
- [50] W. Chen, Y. Liu, H. S. Courtney, M. Bettenga, C. M. Agrawal, J. D. Bumgardner *et al.*, "In vitro anti-bacterial and biological properties of magnetron co-sputtered silver containing hydroxyapatite coating", *Biomaterials* **27** (2006) 5512. <https://www.sciencedirect.com/science/article/abs/pii/S014296120600593X>.
- [51] S. Sharma & M. Kaur, "Antimicrobial activities of rhizobacterial strains of *Pseudomonas* and *Bacillus* strains isolated from rhizosphere soil of Carnation (*Dianthus caryophyllus* Cv. Sunrise)", *Indian Journal of Microbiology* **30** (2010) 229. <https://doi.org/10.1007/s12088-010-0045-6>.
- [52] A. S. Motta, F. Cladera-Olivera & A. Brandelli, "Screening for antimicrobial activity among bacteria isolated from the Amazon basin", *Brazilian Journal of Microbiology* **35** (2004) 307. <https://doi.org/10.1590/S1517-83822004000300007>.
- [53] S. Iqbal & M. Qasim, F. B. Rahman, H. and I. Sajid, "Screening, Characterization and Optimization of antibacterial peptides, produced by *Bacillus safensis* strain MK-12 isolated from waste dump soil KP, Pakistan", *bioRxiv*, 2018, 308205. <https://www.biorxiv.org/content/10.1101/308205v1>.
- [54] K. M. O'Connell, J. T. Hodgkinson, H. F. Sore, M. Welch, G. P. Salmond & D. R. Spring, "Combating multidrug-resistant bacteria: current strategies for the discovery of novel antibacterials", *Angewandte Chemie International Edition* **52** (2013) 10706. <https://onlinelibrary.wiley.com/doi/abs/10.1002/anie.201209979>.
- [55] L. Zhang & C. Sun, "Fengycins, cyclic lipopeptides from marine *Bacillus subtilis* strains, kill the plant-pathogenic fungus *Magnaporthe grisea* by inducing reactive oxygen species production and chromatin condensation", *Applied and Environmental Microbiology* **84** (2018) e00445. <https://pubmed.ncbi.nlm.nih.gov/29980550/>.
- [56] B. Baral & M. R. Mozafari, "Strategic moves of 'superbugs' against available chemical scaffolds: signaling, regulation, and challenges", *ACS Pharmacology & Translational Science* **3** (2020) 373. <https://pubs.acs.org/doi/abs/10.1021/acsp.0c00005>.
- [57] A. Traven & T. Naderer, "Central metabolic interactions of immune cells and microbes: prospects for defeating infections", *EMBO reports* **20** (2019) e47995. <https://www.embopress.org/doi/full/10.15252/embr.201947995>.
- [58] Z. Breijyeh & R. Karaman, "Design and synthesis of novel antimicrobial agents", *Antibiotics* **12** (2023) 628. <https://www.mdpi.com/2079-6382/12/3/628>.